

MAGNETIC SCREENING PROPERTIES OF AN INCOMPRESSIBLE CHIRAL FLUID

Pietro Donatis and Roberto Iengo

*International School for Advanced Studies, I-34014 Trieste, Italy and
Istituto Nazionale di Fisica Nucleare, INFN, Sezione di Trieste, Trieste, Italy*

Abstract

We study the possible penetration of a static magnetic field in an idealized sample of many layers supporting a two dimensional charged chiral quantum fluid, to see whether there is a kind of Meissner effect. This is a non standard problem since the quantum fluid is incompressible having a gap in its spectrum. We find that the system shows an intermediate behaviour between superconducting and non-superconducting fluids, the magnetic field being screened or not depending on its orientation relative to the layers.

1. Introduction.

The study of the physical properties of non-conventional quantum fluids appears to be interesting and promising, even if a precise and detailed comparison with a real existing system is not attempted. In particular, attention has been devoted to possible non-relativistic quantum fluids living in a two-dimensional space, as a framework of models which could be related to layered superconductors (see for instance references [1], [2], [3] in particular for the magnetic properties of realistic layered superconductors). A general approach consists in introducing the problem by means of an effective lagrangian, à la Landau-Ginzburg, representing an universality class which should summarize the relevant degrees of freedom of some underlying microscopic theory not explicitly specified. A particularly interesting class, which is peculiar of two space dimensions, involves somehow a Chern-Simons gauge field, breaking parity and time reversal invariance, describing a non-relativistic quantum fluid which makes a *chiral* distinction between some left- and right-handed behaviour (references [4], [5], [6], [7]).

Here we consider a particular universality class of that kind characterized by the fact that the spectrum of the fluid (before considering it charged and coupled to an electromagnetic field) has a gap. This chiral quantum fluid was presented and extensively studied in ref [7] (it turned out that it is *formally* similar to an effective lagrangian used to describe the Fractional Quantum Hall effect [8], [9], [10], [11]. In our case however, the physical context, the interpretation and the range of the parameters are quite different). It is of course very interesting to study the behaviour of the fluid when it is charged and can carry electromagnetic currents, in order to compare with the behaviour of superconductors. Therefore here we address ourselves more specifically to a rather crucial issue. Due to the fact that this quantum fluid has a gap in the spectrum and it is like an “incompressible” fluid, does it possess the property of screening a magnetic field like a superconductor (Meissner effect) does? Since there is a gap, the standard mechanism similar to the Higgs mechanism in field theory looks impossible. There is no massless Goldstone boson which can provide a “mass term” for the electromagnetic field, thus resulting in a finite penetration length. In fact, in our case, apart from the collective modes representing the overall uniform motion of the fluid, there are not the linearly dispersing compressional modes, which are found in mean field treatments of fluids related to fractional statistics, see in particular [12], [13], [14], [15], [10], (for a possible different treatment, see appendix A of [7] and [16]).

One can make this question more precise for the peculiar quantum fluid we are investigating. Indeed, the reason why there is a gap is due to the fact that it is coupled to a Chern-Simons gauge field $\vec{A}^{CS}$. This field is determined by the constraint that its field strength $\vec{\nabla} \wedge \vec{A}^{CS}$ is proportional to the density fluctuation $\delta\rho$ of the fluid (see section 2 for the formulation of the effective theory). Since the fluid is non-relativistic there is

conservation of the total number of particles, that is $\int \delta\rho = 0$, consistently with the fact that $\vec{\nabla} \wedge \vec{A}^{CS}$ is “exact”, i.e. it is a total derivative. It is because of the CS gauge field that the Goldstone theorem is evaded and that the $U(1)$ phase invariance breaks down, but there is a gap in the spectrum.

Next, when considering the coupling with an electromagnetic field, the sum $(\vec{A}^{CS} + e\vec{A}^{em})$ of the two gauge fields will appear in the covariant derivative. As observed in particular in reference [17], it looks like there remains an unbroken $U(1)$ symmetry, because of a simultaneous gauge transformation of both $\vec{A}^{CS}$ and $\vec{A}^{em}$ which cancels in the sum. A “mass” for the electromagnetic field, and thus the Meissner effect, is recovered in the effective theory of reference [17] by means of a term which separately breaks the gauge invariance for $\vec{A}^{CS}$, due to the condensation of the “spinon pair” component. In our effective theory, instead, we consider a one-component quantum fluid and therefore there is no such a term. The dynamics is therefore quite non standard.

In practice, we can put the question in the following way. The term possibly giving a “mass” to the electromagnetic field, which in the standard case would be proportional to $(e\vec{A}^{em})^2\rho$ (where ρ is the fluid density), is now $(\vec{A}^{CS} + e\vec{A}^{em})^2\rho$. Therefore, there is the possibility that no “mass” and no screening of the magnetic field is obtained, if the configuration

$$\vec{A}^{CS} = -e\vec{A}^{em} \tag{1.1}$$

is energetically favourite. (To avoid possible confusion with the literature on the Hall effect, we remind that in references [8], [9], [10], [11], the relation (1.1) implements the “filling” relation, the CS field being proportional to the total density and the electromagnetic field representing the strong, fixed, space-independent magnetic field of the Hall effect. In our case instead, the CS field strength is proportional to the density *fluctuation*, and the space dependence of the magnetic field is not a priori given, but has to be determined by the variation of the hamiltonian).

We consider here the case of $\vec{A}^{em}$ corresponding to a static magnetic field, and the problem of its penetration in the bulk of a sample made of a stack of many two-dimensional layers, where the quantum fluid lies (see figure 1a-b). In ref [7] we already discussed this problem, finding a different behaviour depending on the orientation of the magnetic field relative to the layers. Namely if the field is orthogonal to the layers’ plane, case of figure 1a, the system behaves like a type II superconductor with a finite penetration length. In the case of figure 1b, when the field is parallel to the layers, instead the penetration length grows with some fractional power of the sample size. Thus in this case there is no Meissner effect, strictly speaking, in agreement with general considerations.

In the study done in reference [7] we have included in the energy computation also the electrostatic effects, arising from the fact that locally $\delta\rho \neq 0$. We imagine of course that the layer where the fluid lies provides a uniform background that neutralizes the charge, so

that the fluid is globally neutral. But assuming that the background charge cannot move, there are locally electrostatic effects where $\delta\rho \neq 0$. These effects arise in particular in the configuration of equation (1.1), due to the fact that $\delta\rho$ is proportional to $\vec{\nabla} \wedge \vec{A}^{CS}$. The resulting additional energy helps in disfavours the configuration (1.1).

However it can well occur that, in more realistic cases, the background-neutralizing-charge is not really fixed and that it can in some way compensate the local excess of electrostatic charge so that the system remains locally neutral. Thus it is important to reconsider the problem.

Here we discuss the central issue of the screening of the magnetic field, by assuming that there are no additional electrostatic terms in the energy. Therefore we examine whether there are intrinsic mechanisms which would energetically disfavour the penetration of the magnetic field.

Our result is that, even in this case, the pattern remains qualitatively the same as said above, namely we find screening, i.e. finite penetration of the magnetic field, in the configuration of figure 1*a* and infinite penetration in the configuration of figure 1*b*, although with some important quantitative differences.

We have summarized in section 3 the main physical reasons for these results, while section 2 summarizes the formulation of the effective theory. Sections 4 and 5 contain the more precise and quantitative analysis.

In conclusion, we find that a chiral quantum fluid, of the universality class described in section 2, besides having other interesting properties discussed in [7], behaves with respect to a static magnetic field as a “quasi-superconductor”, that is in a way which is somewhat intermediate between superconducting and non-superconducting fluids. This could open new perspectives in the scenario of the layered quantum fluids and the possible realistic systems.

Figure 1

2. The effective theory.

The theory which we discuss in this paper is described by the following non-relativistic effective lagrangian density in two space and one time dimensions:

$$\mathcal{L} = i\phi^* \partial_0 \phi - \frac{1}{2m} |\vec{D}\phi|^2 - \frac{\hat{g}}{m} (\delta\rho)^2 \quad (2.1)$$

here $\phi(\vec{x}, t)$ is a non relativistic complex field which plays the rôle of order parameter, related to the density by

$$\rho = |\phi|^2. \quad (2.2)$$

We will comment on the possible values of the dimensionless constant $\hat{g}$ in the next section (in reference [7] we used $g = \frac{\hat{g}}{m}$).

We assume, as it is proper for a non-relativistic theory, a fixed total number of particles N , that is we keep *fixed* $N = \int d^2x \rho$.

$\vec{D}$ is the covariant derivative $\vec{D} = \vec{\nabla} - i\vec{A}^{CS}$ and $\vec{A}^{CS}$ is a Chern-Simons gauge potential related to the matter density fluctuation by the constraint equation:

$$\vec{\nabla} \wedge \vec{A}^{CS} = \frac{2\pi}{k} \delta\rho \quad \delta\rho = \rho - \rho_0 \quad \rho_0 \equiv \langle \rho \rangle. \quad (2.3)$$

k is a dimensionless number which we consider to be of the order of few units. As a consequence of the conservation of the number of particles we have consistently:

$$\int d^2x \vec{\nabla} \wedge \vec{A}^{CS} = 0. \quad (2.4)$$

It is seen that the small deformations have the spectrum

$$E(p) = \sqrt{\mathcal{E}^2 + \frac{1}{4m^2} (\vec{p}^4 + 16\hat{g}\rho_0 \vec{p}^2)} \quad (2.5)$$

where $\mathcal{E} = \frac{2\pi}{mk} \rho_0$ is the gap. The whole spectrum, including the vortex excitations (see also reference [18]), and other relevant properties, in particular the chiral features, have been discussed in ref [7].

Now we are going to study the possible properties of the system of screening an external magnetic field. Since this is essentially a three dimensional phenomenon we suppose to build up a multilayered bulk of many two dimensional thin films separated by a spacing d (see figure 1).

We are interested in stationary situations, and therefore we look for time independent configurations described by the hamiltonian

$$H = \int d^3x \left\{ \frac{1}{2md} |(\vec{\nabla} - ie\vec{A}^{em} - i\vec{A}^{CS})\phi|^2 + \frac{1}{2} (\vec{B}^2 + \vec{E}^2) + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}. \quad (2.6)$$

Here $\vec{B}$ and $\vec{E}$ are the magnetic and electric field, respectively.

3. Qualitative description and summary of the results.

Before going into computational details, let us discuss the essential points of the magnetic field penetration problem, assuming that there is no electrostatic field. In a y -independent configuration, like in figure 1, $A^{CS} = A_y^{CS}(x)$ and $A^{em} = A_y^{em}(x)$ being the only non-vanishing components of the gauge fields, the phase of ϕ can be taken to be a constant, conventionally zero, otherwise it would contribute an additional positive energy (of course it will play a rôle instead in the vortex configuration of section 4.2). Let us look first at the linearized form of the equations coming from the variation of the hamiltonian (dropping higher derivatives terms which are irrelevant for the discussion of zero modes):

$$\begin{aligned} -\partial_x^2 A^{em} + \frac{e\rho_0}{md}(A^{CS} + eA^{em}) &= 0 \\ -\frac{2\hat{g}}{md} \left(\frac{k}{2\pi}\right)^2 \partial_x^2 A^{CS} + \frac{\rho_0}{md}(A^{CS} + eA^{em}) &= 0 \end{aligned} \tag{3.1}$$

One sees immediately that there is in principle a zero mode, corresponding to the configuration of equation (1.1). Thus if (1.1) could be competitive with the standard configuration it would spoil the screening of the external magnetic field. However to understand its relevance, one has to take properly into account the boundary conditions and to see how the allowed modes of A^{CS} can actually implement (1.1). In the following subsections, we will analyze, in a rather qualitative way, the configuration (1.1) in the two geometries of figure 1*a-b*. A more quantitative study is developed in sections 4 and 5. Of course after discussing whether or not the configuration of equation (1.1) is energetically favourite throughout the whole sample, we still have to find the optimum configuration and describe its space dependence. We will discuss it in detail in the following sections using variational methods, to be more general than the linearized approximation.

3.1. Magnetic field orthogonal to the layers' plane.

Let us start discussing the case of figure 1*a*, namely that with $\vec{B}$ orthogonal to the layers' plane. We suppose that the external magnetic field points in the direction of positive z -axis, and penetrates in the bulk in the x direction (see figure 1*a*). We further assume that the matter distribution is uniform in the y and in the z directions and we can choose the electromagnetic and Chern-Simons gauge field pointing in the y direction:

$$\begin{aligned} \vec{A}^{em} &= (0, A^{em}, 0) \\ \vec{A}^{CS} &= (0, A^{CS}, 0) \end{aligned} \tag{3.2}$$

and, in the gauge $\vec{\nabla} \cdot \vec{A}^{em} = \vec{\nabla} \cdot \vec{A}^{CS} = 0$, depending only on x . With these choices the hamiltonian (2.6) becomes ($L_{x,y,z}$ being the sample's sizes in the various directions):

$$\frac{H}{L_y L_z} = \int dx \left\{ \frac{1}{2md} \left(|\partial_x \phi|^2 + |eA^{em} + A^{CS}|^2 \rho \right) + \frac{1}{2} \vec{B}^2 + \frac{1}{2} \vec{E}^2 + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}, \quad (3.3)$$

In this way we have

$$A^{em} = xB \quad \delta\rho = \frac{k}{2\pi} \partial_x A^{CS}. \quad (3.4)$$

From (3.4), using (1.1), we get

$$\delta\rho = -\frac{ke}{2\pi} B \quad (3.5)$$

on every point inside each layer.

Of course, the system is overall electrically neutral, therefore the support over which the quantum fluid lies in each layer acts as the neutralizing background. If we suppose that the neutralizing background charges are fixed, so that the fluctuation of charged matter $\delta\rho$ cannot be locally neutralized, then we have an electric field, inside each layer, given by the equation

$$\vec{\nabla} \cdot \vec{E} = \frac{e}{d} \delta\rho \quad (3.6)$$

here $\frac{\delta\rho}{d}$ is the three dimensional matter density (we remind that since $\delta\rho$ is uniform in the z direction, the z component of $\vec{E}$ is always zero, thus $\vec{E}$ is effectively two dimensional).

Recalling equation (2.3), $\vec{\nabla} \wedge \vec{A}^{CS} = \frac{2\pi}{k} \delta\rho$, we see that $\vec{E}$ and $\vec{A}^{CS}$ are dual two dimensional vectors that is

$$E_i = \frac{ke}{2\pi d} \epsilon_{ij} A_j^{CS} \quad i = x, y. \quad (3.7)$$

This means that the electric field contribution in (3.3) can be rewritten

$$\frac{1}{2} \vec{E}^2 = \frac{k^2 e^2}{4\pi^2 d^2} (\vec{A}^{CS})^2. \quad (3.8)$$

Therefore also when (1.1) holds we recover, through this electrostatic term, a “mass term” for the electromagnetic field $\sim (\vec{A}^{em})^2$. This case has been extensively studied in [7] with the conclusion that the quantum fluid behaves, in this configuration, as a type II superconductor.

Here we are interested in the case when the background charge structure is not so rigid, and we allow the system to neutralize, in some way, the fluctuation $\delta\rho$. Thus, we analyze the behaviour of the system dropping the $\vec{E}^2$ term in the hamiltonian (3.3). We can take into account the energy which is spent by the system for neutralizing the charge fluctuations, while retaining the same form for the effective hamiltonian, by considering

values for the constant $\hat{g}$ of equation (3.3) rather larger than the ones previously considered in reference [7] (there, considerations based on anyon mean field theory suggested $\hat{g} = \pi(1 - \frac{1}{k})$).

We begin by observing that from equation (3.5) we see that the cancellation (1.1) cannot hold everywhere on the two dimensional space. In fact if this were the case we would have

$$\int dx \frac{\delta \rho}{d} = \frac{k}{2\pi d} \int dx \vec{\nabla} \wedge \vec{A}^{CS} = -\frac{ke}{2\pi d} \int dx B = -\frac{ke}{2\pi d} L_x B \quad (3.9)$$

which being different from zero violates the conservation of the number of particles.

Thus, there must be somewhere an additional missing density $\delta \rho_M$. Actually, it will be concentrated on the edge of the sample, otherwise A^{CS} would have a jump and (1.1) would no longer hold afterwards:

$$\delta \rho_M = \frac{ke}{2\pi} B L_x \delta(x - L_x) . \quad (3.10)$$

Notice that

$$\int dx \frac{\delta \rho_M}{d} = \frac{ke}{2\pi d} L_x B \quad (3.11)$$

which exactly compensate (3.9): $\int (\delta \rho + \delta \rho_M) = 0$.

We see that $\delta \rho_M$ is very large since it is proportional to L_x which is macroscopic. Therefore a very large energy comes, for instance, from the term in the hamiltonian which is proportional to $\int (\delta \rho_M)^2$. Thus, we foresee that the configuration of equation (1.1) will be severely energetically disfavoured, and that the quantum fluid will essentially behave, in the configuration of figure 1a, as a standard superconductor. This is confirmed in the detailed analysis of section 4. (It can also be that $\delta \rho_M$, so to speak, disappears because the fluid undergoes locally a kind of phase transition. But if the fluid is stable this too would cost energy, and the conclusion would be the same).

3.2. Magnetic field parallel to the layers' plane.

Let us now turn to the second case in which the external magnetic field is parallel to the layers, that is points in the x direction and penetrates the bulk in the z direction (see figure 1b). Here we suppose uniformity of the matter distribution along y . Again we can choose the gauge fields pointing in the y direction as in (3.2). With these assumptions the gauge electromagnetic field is

$$A^{em} = -zB \quad (3.12)$$

and, if (1.1) holds,

$$\delta \rho = \frac{k}{2\pi} \partial_x A^{CS} = 0 . \quad (3.13)$$

Equation (3.13) holds everywhere but at the border of the sample. In fact, we take formally the usual boundary condition that $\vec{A}^{CS}$ is zero at infinity (that is outside the sample) and from equations (1.1) and (3.13) it is constant inside each layer, i.e. independent of (x, y) :

$$A^{CS} = -eA^{em}(z)\theta(x)\theta(L_x - x) . \quad (3.14)$$

$\theta(x)$ being the usual step function $=0$ for $x < 0$, $=1$ for $x > 0$. Therefore we have:

$$\delta\rho = -\frac{ke}{2\pi}A^{em}(z)[\delta(x) - \delta(L_x - x)] . \quad (3.15)$$

Notice that now not only $\delta\rho=0$ inside each layer, but also $\int dx \frac{\delta\rho}{d}=0$, as the conservation of the total number of particles requires.

Notice also that the total amount of fluid accumulated at each border of every layer is $\int dx \frac{\delta\rho}{d} = \pm \frac{ke}{2\pi}zB$ which remains finite for $L_x \rightarrow \infty$.

We see thus that $\delta\rho$ is not macroscopically large and therefore we expect that its contribution to the energy will not be large.

In reference [7] we have analyzed this configuration including the electrostatic energy which arises when $\delta\rho$ is not neutralized by the background. In this case the electrostatic energy comes from the attraction of the two opposite charges accumulated at the boundaries. This gives a relatively weak effect, because the two boundaries are far apart, and the configuration of equation (1.1) remains energetically favourite.

We will see that is even more so here, when we assume, like in section 3.1, that the background neutralizes also locally $\delta\rho$, and accordingly we forget the electrostatic effects. In section 5. we analyze this case in some detail and confirm that the configuration (1.1) is indeed favourite and find a penetration length l_z , in the z direction, to be of the order of

$$l_z \sim (L_x\delta)^{1/2} \quad (3.16)$$

where L_x is the sample size in the direction of B (we assume $L_y \gtrsim L_x$) and δ is the thickness of the border region where the charge is accumulated (we remind, from reference [7], that including the electrostatic effect one finds $l_z \sim (L_x d^2)^{1/3} \times (\text{logarithmic corrections})$). Thus, in this case it is true that the system behaves differently from an ordinary superconductor, where l_z is finite for $L_x \rightarrow \infty$. Notice however that still $\frac{l_z}{L_x} \rightarrow 0$ for $L_x \rightarrow \infty$.

4. Screening of the magnetic field orthogonal to the layers.

4.1. Edge penetration.

Here we take as a starting point the hamiltonian (3.3) dropping, as said, the electrostatic term:

$$\frac{H}{L_y L_z} = \int dx \left\{ \frac{1}{2md} \left(|\partial_x \phi|^2 + |eA^{em} + A^{CS}|^2 \rho \right) + \frac{1}{2} \vec{B}^2 + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}. \quad (4.1)$$

We imagine that the region where the magnetic field is different from zero is, in absence of the sample, the interval $-s \leq x \leq L_x$. The total flux of the magnetic field is fixed thus:

$$\frac{\Phi}{L_y} \equiv (L_x + s) \cdot B_M = \int_{-s}^{L_x} dx \vec{B} = \text{fixed}. \quad (4.2)$$

Then we redefine the zero of the energy subtracting the constant quantity $-\frac{1}{2} \int dx \vec{B}_M^2$, so that we can rewrite the second term in (3.3) as:

$$\frac{1}{2} \int dx (\vec{B}^2 - \vec{B}_M^2) = \frac{1}{2} \int dx (\vec{B} - \vec{B}_M)^2. \quad (4.3)$$

Therefore the new hamiltonian is;

$$\frac{H}{L_y L_z} = \int dx \left\{ \frac{1}{2md} \left(|\partial_x \phi|^2 + |eA^{em} + A^{CS}|^2 \rho \right) + \frac{1}{2} (\vec{B} - \vec{B}_M)^2 + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}. \quad (4.4)$$

Now we suppose that the sample is placed with an edge at the origin of the x coordinate and that its length in the x -direction is L_x . We imagine that the sample is much smaller than the region where the magnetic field is different from zero, that is $s \gg L_x$.

Here for simplicity we treat the penetration of the magnetic field as if it were uniform, rather than exponentially decaying, and we call l_x the penetration length. Since we have fixed the total value of the magnetic flux we have:

$$\frac{\Phi}{L_y} = (s + l_x) B = (s + L_x) B_M. \quad (4.5)$$

This leads to:

$$B = \frac{s + L_x}{s + l_x} B_M \simeq B_M, \quad (4.6)$$

and to:

$$B - B_M = \left(\frac{s + L_x}{s + l_x} B_M - B_M \right) = \frac{L_x - l_x}{s + l_x} B_M. \quad (4.7)$$

That is:

$$\frac{1}{2} \int_{-s}^{L_x} dx (\vec{B} - \vec{B}_M)^2 \simeq \frac{1}{2} B_M^2 (L_x - l_x) . \quad (4.8)$$

To check the meaningfulness of what we are doing, let us consider the case of the standard superconductor ($\vec{A}^{CS}=0, \rho=\rho_0$) and see what happens to the screening.

The hamiltonian becomes:

$$\frac{H}{L_y L_z} = \frac{1}{2} \int_{-s}^{L_x} dx (\vec{B} - \vec{B}_M)^2 + \frac{e^2 \rho_0}{2md} \int_0^{l_x} dx (A^{em})^2 . \quad (4.9)$$

Notice that since $A^{em}=xB$, we have for the second term in (4.9):

$$\frac{e^2 \rho_0}{2md} \int_0^{l_x} dx B^2 x^2 \simeq \frac{e^2 \rho_0}{2md} B_M^2 \frac{l_x^3}{3} \quad (4.10)$$

that is, using (4.8):

$$\frac{H}{L_y L_z} \simeq \frac{1}{2} B_M^2 \left(L_x - l_x + \frac{e^2 \rho_0}{md} \frac{l_x^3}{3} \right) . \quad (4.11)$$

Minimizing (4.11) with respect to the penetration length l_x we get the standard value

$$l_x = \sqrt{\frac{md}{e^2 \rho_0}} . \quad (4.12)$$

ρ_0/d being the three dimensional mean density. So our assumptions make sense.

Substituting back in equation (4.11) we get the value of the energy of the standard configuration:

$$\frac{H}{L_y L_z} = \frac{1}{2} B_M^2 \left(L_x - \frac{2}{3} l_x \right) . \quad (4.13)$$

4.1.1. Detailed discussion of the configuration $\vec{A}^{CS} = -e\vec{A}^{em}$.

Coming back to the study of our non-standard quantum fluid, as we said there is a possibility of ruining the screening of the magnetic field by means of the cancellation (1.1). If this happens we have:

$$\int dx \frac{\delta \rho}{d} = \frac{k}{2\pi d} \int dx \vec{\nabla} \wedge \vec{A}^{CS} = -\frac{ke}{2\pi d} \int dx B = -\frac{ke}{2\pi d} L_x B_M \quad (4.14)$$

which is far from being zero.

Therefore in order to have the conservation of the number of the particles of the fluid, somewhere there must be some *missing fluid density*, that we will indicate as $\delta\rho_M$, accounting for the mismatch.

Due to the conservation of the number of particles, it follows that $\int dx \delta\rho = 0$, thus it is not possible that (1.1) holds strictly, otherwise $\int dx B \propto \int dx \delta\rho \neq 0$. So we consider a configuration where (1.1) holds as much as it is possible that is everywhere but in a small region, say, of thickness δ . Thus we suppose that $\delta\rho_M$ is concentrated in a small, microscopic, region δ around $x=x_0$, (as we said in section 3.1, $x_0 \sim L_x$), that is:

$$\delta\rho_M = \frac{ke}{2\pi} B_M L_x \frac{1}{\delta\sqrt{\pi}} e^{-\frac{(x-x_0)^2}{\delta^2}} \quad (4.15)$$

in such a way that

$$\int dx \delta\rho_M = \frac{ke}{2\pi} B_M L_x . \quad (4.16)$$

Let us estimate the various contribution to the energy in this configuration.

First:

$$\int dx \frac{1}{2md} |\partial_x \phi|^2 \simeq \frac{1}{2md} \int dx \frac{(\partial_x \delta\rho_M)^2}{4\delta\rho_M} = \frac{1}{4md} \frac{ke}{2\pi} B_M \frac{L_x}{\delta^2} , \quad (4.17)$$

where $\phi = \sqrt{\rho_0 + \delta\rho}$ and thus $\partial_x \phi = \frac{1}{2} \frac{\partial_x \delta\rho_M}{\sqrt{\rho_0 + \delta\rho}} \simeq \frac{1}{2} \frac{\partial_x \delta\rho_M}{\sqrt{\delta\rho_M}}$.

Second:

$$\frac{\hat{g}}{md} \int_0^{L_x} dx \left(-\frac{ke}{2\pi} B_M + \delta\rho_M \right)^2 = \frac{\hat{g}}{md} \left(\frac{ke}{2\pi} \right)^2 B_M^2 L_x \left(\frac{1}{\sqrt{2\pi}} \frac{L_x}{\delta} - 1 \right) . \quad (4.18)$$

We get the total energy

$$\frac{H}{L_y L_z} = \frac{1}{4md} \frac{ke}{2\pi} B_M \frac{L_x}{\delta^2} + \frac{\hat{g}}{md} \left(\frac{ke}{2\pi} \right)^2 B_M^2 L_x \left(\frac{1}{\sqrt{2\pi}} \frac{L_x}{\delta} - 1 \right) . \quad (4.19)$$

We see that the energy gets a contribution proportional to $B_M^2 \frac{L_x^2}{\delta}$. Therefore compared to the energy of the standard configuration, equation (4.13), we see that the configuration implementing the cancellation as in equation (1.1) has an energy which is larger by a macroscopic factor.

Thus, we can disregard the possibility that the configuration (1.1) holds true throughout the whole sample.

4.1.2. A variational analysis.

We study now in some detail the penetration of the magnetic field with a variational approach of the full hamiltonian (4.4). This will allow us to go beyond the linearized approximation, and to take into account possible important non-linear effects. Let us make the following ansatz:

$$\begin{aligned}
B &= B_0 e^{-\lambda x} \quad \Rightarrow \quad A^{em} = -\frac{B_0}{\lambda} e^{-\lambda x} \\
B_M &= B_0 \\
A^{CS} &= \frac{2\pi}{k} \int_0^x dx' \delta\rho(x') = -\alpha \frac{2\pi}{k} \rho_0 x e^{-\mu x} \\
\delta\rho &= \frac{k}{2\pi} \partial_x A^{CS} = -\alpha \rho_0 (1 - \mu x) e^{-\mu x} \\
\rho &= \delta\rho + \rho_0 = \rho_0 \left[1 - \alpha (1 - \mu x) e^{-\mu x} \right].
\end{aligned} \tag{4.20}$$

Note that $\int dx \delta\rho = 0$. The rôle of the parameter α is to leave free the value of $\delta\rho$ at the edge. The goodness of this ansatz can be checked directly on the standard case (4.9) from which we get back the correct value (4.12).

The numerical analysis indicates that $\mu \gg \lambda$. Let us assume it for displaying a somewhat simplified expression, verifying *a posteriori* that $\mu \gg \lambda$ is indeed realized. One gets to the following expression:

$$\frac{md}{L_y L_z} H = \frac{2\pi e B_0 \rho_0^2}{k \mu^2} \frac{1}{\lambda} \alpha + \left(\frac{\pi^2 \rho_0^3}{2k^2 \mu^3} + \frac{1}{4\mu} \hat{g} \rho_0^2 + \mu \rho_0 c(\alpha) \right) \alpha^2 + \frac{e^2 \rho_0 B_0^2}{4} \frac{1}{\lambda^3} - \frac{3md B_0^2}{4} \frac{1}{\lambda}. \tag{4.21}$$

Here

$$c(\alpha) = \frac{1}{8} \int_0^\infty dx \frac{(2-x)^2 e^{-2x}}{1 - \alpha(1-x)e^{-x}} \tag{4.22}$$

is a slowly varying function of α , which for α small tends to a number of order of unit. We will consider it as a constant. Minimizing this expression with respect to α we find

$$\alpha = -\frac{1}{\lambda} Q B_0, \tag{4.23}$$

where $Q = \frac{4\pi e k \mu \rho_0}{2\pi^2 \rho_0^2 + \hat{g} k^2 \mu^2 \rho_0 + 4k^2 \mu^4 c}$. Substituting back in (4.21) and minimizing now with respect to λ we find:

$$l_x \equiv \frac{1}{\lambda} = \sqrt{\frac{md}{e^2 \rho_0}} \sqrt{1 + \frac{16\pi^2 \rho_0^3}{9mdk^2 \mu^4} Q^2 + \frac{4\pi^2 \rho_0}{3ek\mu^2} Q}. \tag{4.24}$$

We see from equation (4.24) that the penetration length approaches the standard one for large $\hat{g}$, for which $Q \rightarrow 0$. We recall again that a large value of $\hat{g}$ is expected because it effectively represents the fact that the system must spend energy to remain electrically neutral when $\delta\rho \neq 0$.

Indeed this results have been confirmed through a numerical minimization with respect to λ and α of the hamiltonian which is obtained from (4.4) using the ansatz (4.20), with $\mu = \sqrt{\rho_0}$ (corresponding to a coherence length of $\sim 10\text{\AA}$), for various values of B_0 . Already for $\hat{g} = 1.5$ we get $l_x \simeq 1.2 \cdot 10^3 \text{\AA}$, and for $\hat{g} = 10$ we get $l_x \simeq 9.8 \cdot 10^2 \text{\AA}$ (the standard value equation (4.12) is $l_x \simeq 8.5 \cdot 10^2 \text{\AA}$). We have also numerically verified that taking μ smaller increases the energy of the configuration, confirming that $\mu \gg \lambda$ as stated above. In fact, for $\mu \rightarrow \infty$ we see that $Q \sim \frac{1}{c} \frac{\pi e \rho_0}{k \mu^3}$ thus $\alpha \rightarrow 0$, and $l_x \rightarrow \sqrt{\frac{e^2 \rho_0}{m d}}$. We see thus from equation (4.21) that the value $\mu \rightarrow \infty$ formally corresponds to the minimal energy. We have taken μ at its physically reasonable maximum value, that is $\mu \sim \sqrt{\rho_0}$.

One can check that α is indeed small for that value, for $\hat{g} \sim 10$ and $B_0 \lesssim 10^3$ gauss one gets $\alpha \lesssim 10^{-2}$ (in all these numerical computations we have taken m to be the mass of the electron, $\sim 250 \text{\AA}^{-1}$, and $\rho_0 = 4 \cdot 10^{-3} \text{\AA}^{-2}$).

We see that the penetration length is independent of the value of B_0 and that α is proportional to $-B_0$ (that is, $\vec{A}^{CS}$ has the sign opposite to $\vec{A}^{em}$, as if the system would like the configuration (1.1) as far as it is possible).

4.2. Vortices.

In this section we study the penetration of the magnetic field from magnetic vortices. The starting point will be the following hamiltonian:

$$\frac{H}{L_z} = \int d^2r \left\{ \frac{1}{2md} \left| \left(\vec{\nabla} - ie\vec{A}^{em} - i\vec{A}^{CS} \right) \phi \right|^2 + \frac{1}{2} \vec{B}^2 + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}. \quad (4.25)$$

We look for solutions of the form:

$$\phi(r, \theta) = f(r) e^{in\theta}, \quad (4.26)$$

n is integer and represent the vorticity.

Substituting (4.26) in (4.25) we get:

$$\frac{H}{L_z} = \int d^2r \left\{ \frac{1}{2md} \left[(\partial_r f)^2 + \frac{1}{r^2} (n - erA_\theta^{em} - rA_\theta^{CS})^2 f^2 \right] + \frac{1}{2} \vec{B}^2 + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}. \quad (4.27)$$

Finiteness of (4.27) requires:

$$\begin{aligned}
f(0) = 0 & \Rightarrow \delta\rho(0) = -\rho_0 \\
\lim_{r \rightarrow \infty} \delta\rho = 0 & \Rightarrow \lim_{r \rightarrow \infty} f = \sqrt{\rho_0} \\
\lim_{r \rightarrow \infty} A_\theta^{em} &= \frac{n}{er} \\
\lim_{r \rightarrow \infty} A_\theta^{CS} &= 0 .
\end{aligned} \tag{4.28}$$

We solve this hamiltonian in a variational way with the following ansatz, which satisfies (4.28):

$$\begin{aligned}
erA_\theta^{em} &= n(1 - e^{-\lambda^2 r^2}) \Rightarrow B = \frac{1}{r} \partial_r (rA_\theta^{em}) = \frac{2n}{e} \lambda^2 e^{-\lambda^2 r^2} \\
rA_\theta^{CS} &= -\frac{\pi}{k} \rho_0 r^2 e^{-\mu^2 r^2} \\
\delta\rho &= \frac{k}{2\pi} \frac{1}{r} \partial_r (rA_\theta^{CS}) = -\rho_0(1 - \mu^2 r^2) e^{-\mu^2 r^2} \\
f^2 &= \rho_0 - \rho_0(1 - \mu^2 r^2) e^{-\mu^2 r^2} .
\end{aligned} \tag{4.29}$$

Notice that $\int_0^\infty d^2r \delta\rho = 0$, therefore there is no missing $\delta\rho_M$. Later on, we will compare this configuration with a configuration where $\vec{A}^{CS} = -e\vec{A}^{em}$ and we will need $\delta\rho_M \neq 0$ like in the discussion of section 4.1.1.

Notice also that in this case, differently from the case treated in the previous section the value of $\delta\rho$ at the origin is fixed to zero by the requirement of finite energy, see equation (4.27), so there is no α parameter.

With this ansatz we get, supposing $\mu \gg \lambda$, to be later verified:

$$\frac{H}{L_z} \simeq \frac{\rho_0 \pi}{2md} \left(\frac{6n\pi\rho_0}{k\mu^2} + n^2 \log \frac{\mu^2}{2\lambda^2} + \frac{2mdn^2\lambda^2}{e^2\rho_0} + \frac{\rho_0\hat{g}}{2\mu^2} \right) . \tag{4.30}$$

Minimizing (4.30) with respect to λ^2 one finds easily:

$$\lambda^2 = \frac{e^2\rho_0}{2md} . \tag{4.31}$$

Therefore we recover the penetration length we had found in [7] studying the case including the electrostatic interaction:

$$l_x = \frac{1}{\lambda} = \sqrt{\frac{2md}{e^2\rho_0}} . \tag{4.32}$$

Notice that this penetration length is independent of $\hat{g}$.

Minimizing (4.30) with respect to μ we find:

$$\mu^2 = \frac{6\pi\rho_0}{nk} + \frac{\hat{g}\rho_0}{2n^2} , \tag{4.33}$$

which corresponds to a coherence length:

$$r_v = \frac{1}{\mu} = \sqrt{\frac{2kn^2}{12n\pi\rho_0 + k\rho_0\hat{g}}} . \quad (4.34)$$

Notice that, as is reasonably expected, the radius r_v of the vortex decreases with increasing $\hat{g}$. All these results have been confirmed through a numerical minimization with respect to λ and μ of the exact hamiltonian with various values of $\hat{g}$. Notice that the results for the vortices agrees fairly well with those for the edge penetration.

In particular for $\hat{g}=2$ we have $r_v \sim 8\text{\AA}$ and $l_x \sim 1167\text{\AA}$, whereas for $\hat{g}=10$ we have $r_v \sim 6\text{\AA}$ and the same value of l_x (corresponding to the value of equation (4.32)). Notice that these results confirm that $\mu \gg \lambda$ as stated above.

Substituting (4.32) and (4.34) in (4.30) we find the energy of the configuration of one vortex:

$$\frac{H}{L_z} \simeq \frac{\rho_0\pi}{2md} \left(\frac{6n\pi\rho_0}{k} r_v^2 + n^2 \log \frac{md}{e^2\rho_0 r_v^2} + n^2 + \frac{\hat{g}\rho_0 r_v^2}{2} \right) . \quad (4.35)$$

4.2.1. The $\vec{A}^{CS} = -e\vec{A}^{em}$ configuration with rotational symmetry.

We now discuss, like in section 4.1.1, the possible cancellation (1.1) in the case of a configuration which has a rotational symmetry, and the flux of B is given, like for the vortices case discussed above. We will compare it with the standard vortex configuration of the previous section. Following the discussion done there, it is unavoidable a missing $\delta\rho_M$. We suppose to have a fixed value of the magnetic flux.

$$\Phi(B) = \int d^2r B = \frac{2\pi}{e} N . \quad (4.36)$$

Therefore

$$\int d^2r \delta\rho = \frac{k}{2\pi} \int d^2r \vec{\nabla} \wedge \vec{A}^{CS} = -\frac{ke}{2\pi} \int d^2r B = -\frac{ke}{2\pi} \pi R_0^2 B = -kN , \quad (4.37)$$

where R_0 is the radius of the sample, supposed to be a disk.

So to have conservation of the number of particles we need a $\delta\rho_M$ of the form:

$$\delta\rho_M = \frac{kN}{2\pi R_0} \frac{1}{\delta\sqrt{\pi}} e^{\frac{(r-R_0)^2}{\delta^2}} , \quad (4.38)$$

where the thickness δ is supposed to be microscopic. Note that

$$\int d^2r \delta\rho_M = kN. \quad (4.39)$$

We take $\delta\rho_M$ concentrated at a macroscopic distance from the vortex core, say at the edge of the sample, that is at $r = R_0$. Otherwise, if it were concentrated at some different point, say at $r = r_0$, then $\vec{A}^{CS}$ would have a jump of $\frac{N}{r}$ at $r = r_0$, due to the fact that $\partial_r(rA^{CS}) = \frac{2\pi}{k}r\delta\rho$, and would no longer cancel eA^{em} for $r > r_0$, contrary to our hypothesis that the cancellation holds throughout the macroscopic size of the sample.

Let us estimate the most relevant contributions to the energy $\frac{H}{L_z}$ coming from the presence of $\delta\rho_M$. We write:

$$\frac{H}{L_z} = \left(\frac{H}{L_z}\right)_1 + \left(\frac{H}{L_z}\right)_2 + \left(\frac{H}{L_z}\right)_3. \quad (4.40)$$

We find first

$$\left(\frac{H}{L_z}\right)_1 = \frac{1}{2md} \int d^2r |D\phi|^2 \simeq \frac{1}{8md} \int d^2r \frac{(\partial_r \delta\rho_M)^2}{\delta\rho_M} = \frac{1}{8md} \frac{kN}{\delta^2}. \quad (4.41)$$

Second:

$$\left(\frac{H}{L_z}\right)_2 = \frac{\hat{g}}{md} \int d^2r (\delta\rho)^2 = \frac{\hat{g}}{md} \left[\sqrt{\pi} \left(\frac{k}{2\pi}\right)^2 \frac{N^2}{R_0\delta} - \frac{k^2}{\pi} \frac{N^2}{R_0^2} \right]. \quad (4.42)$$

Third:

$$\left(\frac{H}{L_z}\right)_3 = \frac{1}{2} \int d^2r B^2 = \frac{2\pi^2}{e^2} \frac{N^2}{\pi R_0^2}. \quad (4.43)$$

We can distinguish two cases.

a) $N = 1$ (or few units). Then the most relevant energy contribution due to $\delta\rho_M$ comes from $\left(\frac{H}{L_z}\right)_1$, since the other two pieces are suppressed, at least, by the factor $\frac{1}{R_0}$, with R_0 macroscopic. This energy (for $\delta^2 \sim \frac{1}{\rho_0}$) is less or equal to the free energy of the standard configuration (4.35). Therefore for a small flux, that is for a very small magnetic field, the configuration where $\vec{A}^{CS}$ cancels $e\vec{A}^{em}$ is possibly favourite.

b) Now we suppose B macroscopic, in other words B is fixed in the macroscopic limit $R_0 \rightarrow \infty$. Therefore from equation (4.37) $N \sim BR_0^2 \rightarrow \infty$. In this case the configuration where $\vec{A}^{CS}$ cancels $e\vec{A}^{em}$ gets the most relevant energy from $\left(\frac{H}{L_z}\right)_2$, namely from the piece proportional to $\frac{N^2}{\delta R_0}$. Therefore, for this configuration

$$\frac{H}{L_z} \simeq \frac{\hat{g}}{4md} \sqrt{\pi} \left(\frac{ke}{2\pi}\right)^2 B^2 \frac{R_0^3}{\delta} + \text{less important}. \quad (4.44)$$

We have to compare it with the energy of the standard vortex configuration, equation (4.35) multiplied by N , that is:

$$\frac{H}{L_z} \simeq c \frac{\rho_0 \pi}{2md} B R_0^2, \quad (4.45)$$

where c is of the order of $\rho_0 r_v^2$, i.e. a finite number. Since $\frac{R_0}{\delta} \rightarrow \infty$, clearly the standard configuration, or also a configuration of many standard vortices, is energetically favourite.

5. Weak screening of the magnetic field parallel to the layers.

In this chapter we put our attention to the configuration in which the field $\vec{B}$ is parallel to the layers' plane (see figure 1b) and study the screening effects in absence of the electrostatic interaction, as discussed in the previous chapter.

With considerations very similar to those made at the beginning of section 4.1 we get, in this configuration, to the hamiltonian:

$$\frac{H}{L_y} = \int dx dz \left\{ \frac{1}{2md} \left[|\partial_x \phi|^2 + c_J |\partial_z \phi|^2 \right] + \frac{1}{2md} |eA^{em} + A^{CS}|^2 \rho + \frac{1}{2} (B - B_M)^2 + \frac{\hat{g}}{md} (\delta\rho)^2 \right\}, \quad (5.1)$$

here c_J is a constant accounting for the Josephson coupling between the layers.

We study the possible screening starting from the configuration in which we have the cancellation (1.1) inside the sample, that is:

$$A^{CS}(x, z) = -eA^{em}(z) [\theta(x)\theta(L_x - x)] . \quad (5.2)$$

This yields (see section 3.2):

$$\delta\rho(x, z) = \frac{k}{2\pi} \partial_x A^{CS} = -\frac{ke}{2\pi} A^{em} [\delta(x) - \delta(L_x - x)] . \quad (5.3)$$

Here, as in chapter 4, we suppose that the fluid density is confined in a microscopic region of thickness δ so we approximate the δ -functions with

$$\delta(x) \simeq \frac{1}{\sqrt{\pi}\delta} e^{-\frac{x^2}{\delta^2}} . \quad (5.4)$$

Therefore we have:

$$\delta\rho(x, z) = -\frac{ke}{2\pi} \frac{1}{\sqrt{\pi}\delta} \left[e^{-\frac{x^2}{\delta^2}} - e^{-\frac{(L_x - x)^2}{\delta^2}} \right] A^{em}(z) . \quad (5.5)$$

Keeping this configuration, let us assume that the magnetic field penetrates in the z direction within a length l_z and let us estimate it. We begin by estimating the various contributions to the energy in this configuration disregarding the terms proportional to $e^{-\frac{L_x^2}{2\delta^2}} \simeq 0$. First:

$$\frac{1}{2md} \int dx dz |\partial_x \phi|^2 = \frac{1}{4md\rho_0} \frac{1}{\sqrt{2\pi}} \left(\frac{ke}{2\pi} \right)^2 \frac{1}{\delta^3} \int_0^{l_z} dz \left(A^{em}(z) \right)^2 , \quad (5.6)$$

where we used the fact that $\partial_x \phi = \frac{1}{2} \frac{\partial_x(\delta\rho)}{\sqrt{\delta\rho + \rho_0}} \simeq \frac{1}{2\sqrt{\rho_0}} \partial_x(\delta\rho)$.

Second:

$$\frac{c_J}{2md} \int dx dz |\partial_z \phi|^2 = \frac{c_J}{8md\rho_0} \frac{1}{\sqrt{2\pi}} \left(\frac{ke}{2\pi} \right)^2 \frac{1}{\delta} \int_0^{l_z} dz \left(\partial_z A^{em}(z) \right)^2 . \quad (5.7)$$

Third:

$$\frac{\hat{g}}{md} \int dx dz (\delta\rho)^2 = \frac{\hat{g}}{md} \frac{1}{\sqrt{2\pi}} \left(\frac{ke}{2\pi} \right)^2 \frac{1}{\delta} \int_0^{l_z} dz \left(A^{em}(z) \right)^2 . \quad (5.8)$$

Fourth:

$$\frac{1}{2} \int dx dz (B - B_M)^2 = \frac{1}{2} B_M^2 L_x (L_z - l_z) , \quad (5.9)$$

(here to get (5.9) we have used arguments similar to those that led to (4.8)).

Now putting it all together, and using the fact that $A^{em}(z) = -zB_M$, we arrive at:

$$\begin{aligned} \frac{H}{L_y} = & \frac{1}{md} \frac{1}{\sqrt{2\pi}} \left(\frac{ke}{2\pi} \right)^2 \frac{1}{\delta} \left(\frac{1}{4\rho_0\delta^2} + \hat{g} \right) B_M^2 \frac{l_z^3}{3} + \frac{c_J}{8md\rho_0} \frac{1}{\sqrt{2\pi}} \left(\frac{ke}{2\pi} \right)^2 \frac{1}{\delta} B_M^2 l_z + \\ & + \frac{1}{2} B_M^2 L_x (L_z - l_z) . \end{aligned} \quad (5.10)$$

Minimizing equation (5.10) with respect to l_z we find:

$$l_z \simeq \sqrt{\frac{L_x}{\Omega}} , \quad (5.11)$$

where $\Omega = \frac{2}{md} \frac{1}{\sqrt{2\pi}} \left(\frac{ke}{2\pi} \right)^2 \frac{1}{\delta} \left(\frac{1}{4\rho_0\delta^2} + \hat{g} \right)$.

This means that when the magnetic field is parallel we do not have screening in the usual sense because $l_z \rightarrow \infty$ for $L_x \rightarrow \infty$. But we have a “quasi screening” in the sense that $\frac{l_z}{L_x} \rightarrow 0$.

Substituting this result back in (5.10), we find the energy of the configuration:

$$\frac{H}{L_y} = \frac{1}{2} B_M^2 L_x L_z - \frac{1}{3} B_M^2 (L_x)^{3/2} (\Omega)^{-1/2} . \quad (5.12)$$

Let us compare this result to the energy of the standard configuration. That is, let us take $A^{CS} = 0$ and $\phi = \sqrt{\rho_0}$; then the hamiltonian becomes:

$$\frac{H}{L_y} = \frac{1}{2} B_M^2 L_x \left(\frac{e^2 \rho_0}{3md} l_z^3 + L_z - l_z \right) , \quad (5.13)$$

which is minimal for the standard value (4.12) of the penetration length. The value of this minimal energy is:

$$\frac{H}{L_y} = \frac{1}{2} B_M^2 L_x \left(L_z - \frac{2}{3} l_z \right) . \quad (5.14)$$

Comparing this equation with equation (5.12) we see that (5.14) is always greater than (5.12) for L_x macroscopic. So the configuration (1.1), in the geometry as in fig 1b is always favourite.

Acknowledgements. We would like to thank Arturo Tagliacozzo for useful comments on our previous work, which have stimulated the present paper.

References.

- [1] W.E. LAWRENCE, S. DONIACH, in “Proceedings of the XII International Conference on Low Temperature Physics” (E. Kanda, Ed.), p.361, Academic Press, Tokyo, 1971.
- [2] L.N. BULAEVSKII, *Int. J. Mod. Phys.* **B4** (1990), 1849.
- [3] J.R. CLEM, *Phys. Rev.* **B43** (1991), 7837.
- [4] J. MARCH-RUSSELL, F. WILCZEK, *Phys. Rev. Lett.* **61** (1988), 2066.
- [5] B.I. HALPERIN, J. MARCH-RUSSELL, F. WILCZEK, *Phys. Rev.* **B40** (1989), 8726.
- [6] X.G. WEN, F. WILCZEK, A. ZEE, *Phys. Rev.* **B39** (1989), 11413.
- [7] P. DONATIS, R. IENGO, *Nucl. Phys.* **B[FS]415** (1994), 630.
- [8] S.C. ZHANG, T.H. HANSSON, S. KIVELSON, *Phys. Rev. Lett.* **62** (1989), 82.
- [9] D.H. LEE, S.C. ZHANG, *Phys. Rev. Lett.* **66** (1991), 1220.
- [10] D.H. LEE, M.P.A. FISHER, *Int. J. Mod. Phys.* **B5** (1991), 2675.
- [11] S.C. ZHANG, *Int. J. Mod. Phys.* **B6** (1992), 25.
- [12] A.L. FETTER, C.B. HANNA, R.B. LAUGHLIN, *Phys. Rev.* **B39** (1989), 9679.
- [13] Y.H. CHEN, F. WILCZEK, E.WITTEN, B.I. HALPERIN, *Int. J. Mod. Phys.* **B3** (1989), 1001.
- [14] A.L. FETTER, C.B. HANNA, R.B. LAUGHLIN, *Phys. Rev.* **B40** (1989), 8745.
- [15] X.G. WEN, A. ZEE, *Phys. Rev.* **B41** (1990), 240.
- [16] R. IENGO, K. LECHNER, *Nucl. Phys.* **B384** (1992), 541.
- [17] X.G. WEN, A.ZEE, *Phys. Rev. Lett.* **62** (1989), 2873.
- [18] P. DONATIS, R. IENGO, *Phys. Lett.* **B320** (1994), 64.